

Lepton asymmetry and primordial nucleosynthesis in the era of precision cosmology

Pasquale D. Serpico and Georg G. Raffelt

Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), Föhringer Ring 6, 80805 München, Germany

(Dated: February 2, 2008)

We calculate and display the primordial light-element abundances as a function of a neutrino degeneracy parameter ξ common to all flavors. It is the only unknown parameter characterizing the thermal medium at the primordial nucleosynthesis epoch. The observed primordial helium abundance Y_p is the most sensitive cosmic “leptometer.” Adopting the conservative Y_p error analysis of Olive and Skillman implies $-0.04 \lesssim \xi \lesssim 0.07$ whereas the errors stated by Izotov and Thuan imply $\xi = 0.0245 \pm 0.0092$ (1σ). Improved determinations of the baryon abundance have no significant impact on this situation. A determination of Y_p that reliably distinguishes between a vanishing or nonvanishing ξ is a crucial test of the cosmological standard assumption that sphaleron effects equilibrate the cosmic lepton and baryon asymmetries.

PACS numbers: 26.35.+c, 14.60.Lm, 98.80.Es

Preprint MPP-2005-54

I. INTRODUCTION

The successful calculation of the primordial light-element abundances in the framework of the standard theory of big-bang nucleosynthesis (BBN) [1, 2, 3, 4] is among the main pillars supporting our modern understanding of the universe. Apart from the cross sections that are needed for the network of nuclear reactions, these predictions depend on the baryon-to-photon ratio η_B , on the radiation density at the BBN epoch that is traditionally parametrized by the equivalent number of neutrino flavors N_{eff} , on the electron-neutrino degeneracy parameter $\xi = \mu_{\nu_e}/T_{\nu_e}$, and on the degeneracy parameters of the other neutrino flavors. Originally all of these parameters had to be fitted such as to reproduce the observed light-element abundances.

Meanwhile the number of neutrino flavors has been fixed by the Z^0 decay width [4]. More recently, the baryon abundance has been determined by the cosmic microwave background (CMB) anisotropies to be [5]

$$\eta_B = \frac{n_B - n_{\bar{B}}}{n_\gamma} = 6.14 \times 10^{-10} (1.00 \pm 0.04). \quad (1)$$

Finally, the measured neutrino mixing parameters imply that neutrinos reach approximate chemical equilibrium before the BBN epoch so that all neutrino chemical potentials can be taken to be equal, i.e. they are all characterized by the same degeneracy parameter ξ that applies to ν_e [6, 7, 8].

Of course, one may still assume new low-energy physics such as sizeable active-sterile neutrino mixings, the existence of new particles such as axions that could contribute to the primordial radiation density, a time-varying Newton constant, and a plethora of other non-standard options that would affect BBN [9]. Barring these more exotic possibilities, the single remaining BBN parameter that is not otherwise fixed is ξ . In the light of these developments we re-examine BBN under the assumption that ξ is the only free input parameter.

II. BBN WITH A LEPTON ASYMMETRY

Over the years, BBN with a lepton asymmetry has been studied by many authors [1, 10, 11, 12, 13, 14, 15, 16]. To achieve approximate agreement between the observed and predicted light-element abundances a possible lepton asymmetry must be small so that one may well constrain a new analysis, say, to the range $|\xi| < 0.5$. For such small ξ values the most important impact on BBN is a shift of the beta equilibrium between protons and neutrons. Subleading effects include a modification of the radiation density,

$$\Delta N_{\text{eff}} = 3 \left[\frac{30}{7} \left(\frac{\xi}{\pi} \right)^2 + \frac{15}{7} \left(\frac{\xi}{\pi} \right)^4 \right]. \quad (2)$$

Moreover, the neutrino decoupling temperature is higher than in the standard case [10, 11] so that in principle one could get a non-standard $T_\nu(T)$ evolution, but such effects are completely negligible for our case. A non-zero ξ slightly modifies the partial neutrino reheating following the e^+e^- annihilation [12], again a completely negligible effect for our ξ range of interest.

To include the modified $n \leftrightarrow p$ weak rates we adopt a perturbative approach for the modified neutrino distribution functions, similar to the one adopted by us in a different context [17]. Therefore, the additional terms for the rates can be obtained by integrating the factors

$$\delta f_\nu(E, \xi) = \sum_{n=1}^{\infty} \frac{\partial f_\nu(E, \xi)}{\partial \xi} \bigg|_{\xi=0} \frac{\xi^n}{n!}. \quad (3)$$

Corrections up to fourth order were included into the code of Ref. [18], thus reducing the error of the truncation on the nuclide yields to the fourth significant digit for $|\xi| \leq 0.1$, and to the third for $|\xi| \leq 0.5$. This also sets the level of the required fitting accuracy of the auxiliary functions. The corrections were calculated as relative changes with respect to the $\xi = 0$ Born rates so that in the $\xi \rightarrow 0$ limit we recover the standard $n \leftrightarrow p$ rates with finite mass, QED radiative and thermodynamic corrections, and partial neutrino reheating [18]. For $\xi \neq 0$




FIG. 1: Light-element abundances as a function of the neutrino degeneracy parameter. The top panel shows the primordial ${}^4\text{He}$ mass fraction Y_p whereas the other panels show the ${}^2\text{H}$, ${}^3\text{He}$, and ${}^7\text{Li}$ number fractions relative to hydrogen. The gray 1σ error bands include the uncertainty of the WMAP determination of the baryon abundances as well as the uncertainties from the nuclear cross sections [18]. We also show the observed abundances as described in the text.

these sub-leading effects are rescaled without introducing significant errors. We have checked the reliability of our approach by comparing with the code of Ref. [13] where a fully numerical and non-perturbative method is used.

In Fig. 1 we show our predictions for the primordial light-element abundances as a function of the neutrino degeneracy parameter ξ , taken to be equal for all flavors. The gray band is the 1σ predicted range. We used the




FIG. 2: A blowup of the first panel of Fig. 1, showing the 1σ range for the primordial helium mass fraction. The small box represents the 1σ observational error quoted by Izotov and Thuan [19] whereas the large box represent the maximum systematic error quoted by Olive and Skillman [20].

nuclear reactions and uncertainties adopted in Ref. [18]. However, except for lithium the uncertainty is dominated by the error of the Wilkinson microwave anisotropy probe (WMAP)-implied [5] baryon abundance of Eq. (1). In Fig. 2 we show a blowup of the top panel, i.e. the predicted helium abundance.

We also show observations of the primordial abundances. Beginning with ${}^7\text{Li}$ (bottom panel of Fig. 1) we show the allowed range $\log_{10}(X_7) + 12 = 2.0\text{--}2.3$, where the uncertainty on T_{eff} has been included [21]. In the second to last panel we show Bania et al.'s possible upper limit on the ${}^3\text{He}$ abundance of 1.3×10^{-5} [22] (at 1σ). The status and the interpretation of these observations is quite controversial, especially for ${}^7\text{Li}$; see Ref. [18] for further discussions and references. It is evident, however, that the ${}^3\text{He}$ upper limit is compatible with any realistic ξ value whereas ${}^7\text{Li}$ is always incompatible. Rather than dismissing standard BBN as the correct theory for the light-element synthesis we follow the usual practice of dismissing the ${}^7\text{Li}$ observations as too uncertain to be useful. Putting this more positively, it is remarkable how close the ${}^7\text{Li}$ abundances observed on the Spite plateau in metal poor stars come to the BBN prediction.

Turning to deuterium (second panel in Fig. 1), we note that a new detection was reported, giving $X_2 \equiv {}^2\text{H}/\text{H} = 1.60^{+0.25}_{-0.30} \times 10^{-5}$ [23], significantly lower than the previous value towards Q1243+3047 of $2.42^{+0.35}_{-0.25} \times 10^{-5}$ [24]. By adopting the conservative approach to symmetrize the errors to the higher value, and the wider allowed range $(1.57\text{--}4.00) \times 10^{-5}$ for the most uncertain determination towards Q0347-3819, one gets 2.42 ± 0.27 (outer error box in Fig. 1) for the average value of the seven positive detections so far obtained in quasar Lyman absorption systems. As previously noted [24], since χ^2 per degree of freedom is approximately 4.4, even with these gener-

ous error estimates, underestimated systematics may be present.

Still, the observed primordial deuterium abundance looks perfectly compatible with the standard BBN predictions in the absence of a lepton asymmetry. Neglecting possible systematics one finds a lower limit $\xi > -0.2$ at 1σ , but no useful upper limit obtains. This remains true even if the precision of the deuterium observations improve by a factor of two (inner box in Fig. 1).

From Figs. 1 and 2 it is obvious that ${}^4\text{He}$ is the best probe of ξ . As the dependence on the baryon abundance η_B is only logarithmic, the new WMAP data do not significantly improve the constraint. Therefore, the observed range for Y_p is the only truly significant measure for a possible lepton asymmetry.

Recently new data were published by Izotov and Thuan [19], giving $Y_p = 0.2421 \pm 0.0021$ (inner error box in Figs. 1 and 2). Slightly higher values arise for differently chosen samples and/or for other metallicity regressions. For the purpose of illustration we have performed a likelihood analysis of the BBN prediction for $Y_p(\eta_B, \xi)$ using the η_B prior of Eq. (1) and Izotov and Thuan's Y_p with their quoted statistical error. The result suggests a positive value $\xi = 0.0245 \pm 0.0092$ (1σ), i.e. a hint for $\xi \neq 0$ at 2.7σ . This exercise illustrates the possible sensitivity that present determinations of ${}^4\text{He}$ could reach if the systematic uncertainties were fully understood.

In an independent analysis Olive and Skillman [20] find $Y_p = 0.249 \pm 0.009$. In an attempt to quantify systematic effects, they suggest as the most conservative range $0.232 \leq Y_p \leq 0.258$ (outer box in Figs. 1 and 2), corresponding to $-0.044 \lesssim \xi \lesssim 0.070$ for the 1σ range of η_B or $-0.046 \lesssim \xi \lesssim 0.072$ for the 2σ range.

To compare ξ with η_B on the same footing we note that the cosmic abundance of a lepton flavor α is given in terms of the corresponding degeneracy parameter as

$$\eta_\alpha = \frac{n_\alpha - n_{\bar{\alpha}}}{n_\gamma} = \frac{1}{12\zeta(3)} \left(\frac{T_\alpha}{T_\gamma} \right)^3 (\pi^2 \xi_\alpha + \xi_\alpha^3). \quad (4)$$

Charge neutrality implies that η_e is of the same order as η_B . A possible large lepton asymmetry thus resides in neutrinos,

$$\eta_L \approx \sum_\nu \eta_\nu \approx 3 \times 0.249 \xi, \quad (5)$$

where $(T_\nu/T_\gamma)^3 = 4/11$ was assumed. Note that this factor is missing in Eq. (1) of Ref. [14] and Eq. (10) of Ref. [16], that limits their validity to temperatures larger than the electron mass.

III. DISCUSSION AND SUMMARY

We have argued that after the experimental determination of the neutrino mixing parameters and after the WMAP determination of the baryon abundance, of all parameters that characterize the cosmic thermal heat

bath at the BBN epoch only the lepton asymmetry, i.e. a neutrino degeneracy parameter ξ common to all flavors, remains undetermined and has to be fixed by the observed light-element abundances. For the first time we have produced a meaningful plot of the predicted light-element abundances as a function of ξ . One motivation for writing this short note was to present for the first time this plot as an alternative to the still common but outdated depiction of the light-element abundances as a function of η_B .

The errors of the predictions in our plot are typically dominated by the WMAP determination of the baryon fraction η_B and thus will significantly improve with forthcoming CMB missions.

The deuterium abundance is very sensitive to η_B so that deuterium is the best baryometer of all the light elements. On the other hand, this element does not respond much to a non-zero lepton asymmetry. The opposite applies to helium which is by far the most sensitive leptometer whereas it is virtually insensitive to future improvements of η_B determinations.

The usual attitude towards the possibility of a large cosmic lepton asymmetry is that sphaleron effects before electroweak symmetry breaking equilibrate the cosmic lepton and baryon asymmetries to within a factor of order unity. In the standard model one finds that $\eta_L = -\frac{51}{28}\eta_B$. From this perspective, BBN is a parameter-free theory and the cosmic helium abundance is fixed by the $\xi = 0$ value shown in Fig. 2, i.e. $Y_p = 0.2479 \pm 0.0005$ (1σ). In this scenario it remains to understand the systematic errors in the spectroscopic Y_p determinations.

However, we think that this argument should be turned around. Sphaleron effects are a crucial ingredient in most baryogenesis scenarios [25], including leptogenesis [26]. No experimental evidence for or against these effects exists. Therefore, a possible indication for a nonzero ξ by the observed helium abundance presents an approach for a possible falsification of the usual assumptions about baryon-lepton equilibration.

In principle, other cosmological probes of a non-zero ξ exist in the form of the CMB anisotropies and large-scale structure (LSS) power spectra [27, 28]. The former essentially feels a boost in the amplitude of the first peak and a shift towards higher multipoles, effects that are degenerate with $N_{\text{eff}} > 3$. The latter suffers a suppression very similar to the effect of a non-zero m_ν . However, the current CMB and LSS data are only sensitive to $\xi = \mathcal{O}(1)$ [29].

A step further may be provided by future CMB limits on ΔN_{eff} , possibly helped by accurate polarization maps [30]. Assuming Kinney and Riotto's estimate [27] of the capability of PLANCK to detect one degeneracy parameter of order 0.5, we can easily translate it into a marginal sensitivity of $\xi \simeq 0.3$ that is common to all three flavors. However, unless one is able to break the degeneracies with other cosmological parameters, such a level of accuracy is too optimistic.

To reach a sensitivity comparable with BBN, one has

to wait for more ambitious methods studying the gravitational lensing distortions on both the temperature and polarization maps of the CMB [31]. Emphasis is usually put on the sensitivity to neutrino masses, but these measurements are useful to constrain the relic neutrino asymmetry as well, especially if complementary information would be provided by improved LSS maps and direct experimental data on the neutrino masses. Incidentally, such an accurate satellite mission would be sensitive to Y_p at the recombination epoch at a level of $\delta Y_p \lesssim 0.005$ [31], thus comparable or better than present astrophysical spectroscopic determinations, with a positive impact on the BBN constraint on ξ . In any event, BBN is the only probe sensitive to the sign of ξ .

In summary, now that the cosmic baryon abundance has been extremely well determined by CMB observa-

tions and now that much about neutrino properties has been learnt by experiments, the role of BBN as a baryometer has shifted to that of the best available cosmic leptometer. Therefore, a more reliable Y_p determination is of much greater fundamental interest than the next round of more precise CMB baryon determinations.

ACKNOWLEDGMENTS

We acknowledge partial support by the Deutsche Forschungsgemeinschaft under Grant No. SFB-375 and by the European Union under the Ilias project, contract No. RII3-CT-2004-506222.

-
- [1] R. V. Wagoner, W. A. Fowler and F. Hoyle, "On the synthesis of elements at very high temperatures," *Astrophys. J.* **148**, 3 (1967).
 - [2] K. A. Olive, G. Steigman and T. P. Walker, "Primordial nucleosynthesis: Theory and observations," *Phys. Rept.* **333**, 389 (2000) [astro-ph/9905320].
 - [3] S. Burles, K. M. Nollett and M. S. Turner, "Big-bang nucleosynthesis predictions for precision cosmology," *Astrophys. J.* **552**, L1 (2001) [astro-ph/0010171].
 - [4] S. Eidelman *et al.* (Particle Data Group), "Review of particle physics," *Phys. Lett. B* **592**, 1 (2004).
 - [5] D. N. Spergel *et al.* (WMAP Collaboration), "First Year Wilkinson Microwave Anisotropy Probe (WMAP) observations: Determination of cosmological parameters," *Astrophys. J. Suppl.* **148**, 175 (2003) [astro-ph/0302209].
 - [6] A. D. Dolgov, S. H. Hansen, S. Pastor, S. T. Petcov, G. G. Raffelt and D. V. Semikoz, "Cosmological bounds on neutrino degeneracy improved by flavor oscillations," *Nucl. Phys. B* **632**, 363 (2002) [hep-ph/0201287].
 - [7] Y. Y. Y. Wong, "Analytical treatment of neutrino asymmetry equilibration from flavour oscillations in the early universe," *Phys. Rev. D* **66**, 025015 (2002) [hep-ph/0203180].
 - [8] K. N. Abazajian, J. F. Beacom and N. F. Bell, "Stringent constraints on cosmological neutrino antineutrino asymmetries from synchronized flavor transformation," *Phys. Rev. D* **66** (2002) 013008 [astro-ph/0203442].
 - [9] S. Sarkar, "Big bang nucleosynthesis and physics beyond the standard model," *Rept. Prog. Phys.* **59**, 1493 (1996) [hep-ph/9602260].
 - [10] K. Freese, E. W. Kolb and M. S. Turner, "Massive, degenerate neutrinos and cosmology," *Phys. Rev. D* **27**, 1689 (1983).
 - [11] H. S. Kang and G. Steigman, "Cosmological constraints on neutrino degeneracy," *Nucl. Phys. B* **372**, 494 (1992).
 - [12] S. Esposito, G. Miele, S. Pastor, M. Peloso and O. Pisanti, "Non equilibrium spectra of degenerate relic neutrinos," *Nucl. Phys. B* **590**, 539 (2000) [astro-ph/0005573].
 - [13] S. Esposito, G. Mangano, G. Miele and O. Pisanti, "The standard and degenerate primordial nucleosynthesis versus recent experimental data," *JHEP* **0009**, 038 (2000) [astro-ph/0005571].
 - [14] V. Barger, J. P. Kneller, P. Langacker, D. Marfatia and G. Steigman, "Hiding relativistic degrees of freedom in the early universe," *Phys. Lett. B* **569**, 123 (2003) [hep-ph/0306061].
 - [15] A. Cuoco, F. Iocco, G. Mangano, G. Miele, O. Pisanti and P. D. Serpico, "Present status of primordial nucleosynthesis after WMAP: Results from a new BBN code," *Int. J. Mod. Phys. A* **19**, 4431 (2004) [astro-ph/0307213].
 - [16] J. P. Kneller and G. Steigman, "BBN For Pedestrians," *New J. Phys.* **6**, 117 (2004) [arXiv:astro-ph/0406320].
 - [17] P. D. Serpico and G. G. Raffelt, "MeV-mass dark matter and primordial nucleosynthesis," *Phys. Rev. D* **70** (2004) 043526 [astro-ph/0403417].
 - [18] P. D. Serpico, S. Esposito, F. Iocco, G. Mangano, G. Miele and O. Pisanti, "Nuclear reaction network for primordial nucleosynthesis: A detailed analysis of rates, uncertainties and light nuclei yields," *JCAP* **0412**, 010 (2004) [astro-ph/0408076].
 - [19] Y. I. Izotov and T. X. Thuan, "Systematic effects and a new determination of the primordial abundance of ^4He and dY/dZ from observations of blue compact galaxies," *Astrophys. J.* **602**, 200 (2004) [astro-ph/0310421].
 - [20] K. A. Olive and E. D. Skillman, "A realistic determination of the error on the primordial helium abundance: Steps toward nonparametric nebular helium abundances," *Astrophys. J.* **617**, 29 (2004) [astro-ph/0405588].
 - [21] D. L. Lambert, "Lithium in very metal-poor dwarf stars — problems for standard big bang nucleosynthesis?," *AIP Conf. Proc.* **743** (2005) 206 [astro-ph/0410418].
 - [22] T. M. Bania, R. T. Rood and D. S. Balser, "The cosmological density of baryons from observations of $^3\text{He}^+$ in the Milky Way," *Nature* **415**, 54 (2002).
 - [23] N. H. M. Crighton, J. K. Webb, A. Ortiz-Gill and A. Fernandez-Soto, "D/H in a new Lyman limit quasar absorption system at $z = 3.256$ towards PKS1937-1009," *Mon. Not. Roy. Astron. Soc.* **355**, 1042 (2004) [astro-ph/0403512].
 - [24] D. Kirkman, D. Tytler, N. Suzuki, J. M. O'Meara and D. Lubin, "The cosmological baryon density from the deuterium to hydrogen ratio towards QSO absorp-

- tion systems: D/H towards Q1243+3047,” *Astrophys. J. Suppl.* **149**, 1 (2003) [astro-ph/0302006].
- [25] M. Dine and A. Kusenko, “The origin of the matter-antimatter asymmetry,” *Rev. Mod. Phys.* **76**, 1 (2004) [hep-ph/0303065].
 - [26] W. Buchmüller, R. D. Peccei and T. Yanagida, “Leptogenesis as the origin of matter,” hep-ph/0502169.
 - [27] W. H. Kinney and A. Riotto, “Measuring the cosmological lepton asymmetry through the CMB anisotropy,” *Phys. Rev. Lett.* **83**, 3366 (1999) [hep-ph/9903459].
 - [28] J. Lesgourgues and S. Pastor, “Cosmological implications of a relic neutrino asymmetry,” *Phys. Rev. D* **60**, 103521 (1999) [hep-ph/9904411].
 - [29] S. Hannestad, “Neutrino masses and the number of neutrino species from WMAP and 2dFGRS,” *JCAP* **0305**, 004 (2003) [astro-ph/0303076].
 - [30] R. E. Lopez, S. Dodelson, A. Heckler and M. S. Turner, “Precision detection of the cosmic neutrino background,” *Phys. Rev. Lett.* **82**, 3952 (1999) [astro-ph/9803095].
 - [31] M. Kaplinghat, L. Knox and Y. S. Song, “Determining neutrino mass from the CMB alone,” *Phys. Rev. Lett.* **91**, 241301 (2003) [astro-ph/0303344].